



# Impacts of Unmanaged Year-long Grazing




Rangelands are ecosystems that have adapted to withstand such disturbances as drought, flood, fire, and grazing. All disturbances affect plants to some extent, either directly or indirectly, depending on the **timing**, **intensity**, and **frequency** of the disturbance. Generally, the more diverse the vegetation, the better rangeland can withstand disturbance.

Rangeland plants provide nutrients—proteins, starches, and sugars—to grazing and browsing livestock and wildlife. These nutrients, or plant foods, are produced by photosynthesis. Because photosynthesis occurs only in green plant tissue and mostly in the leaves, a plant becomes less able to produce food, at least temporarily, when its leaves are removed (defoliation) by grazing and browsing animals.

Products of photosynthesis are just as important to plants as they are to animals. Like all other living things, plants need food to survive and grow. The food that plants make for themselves through photosynthesis is used for major plant functions such as surviving dormancy, growing new roots, growing new leaves in the spring, and replacing leaves lost to grazing or browsing.<sup>1</sup>

**Failure to manage grazing can lead to overuse of plants. As the graphic below shows, when too many leaves are removed from the plant, the growth and vigor of plant root systems slows or stops. This can reduce carbon input into the soil, result in less resilience to drought, and even grass mortality, increased soil erosion, and soil that is less able to capture water.**

“Short roots can’t hold the soil in place, let alone do their job of feeding soil microorganisms, creating a sponge to hold water, and pulling carbon from the air deep, deep into the soil where it can be sequestered. It’s that depth that makes the difference between carbon that “breathes” back and forth between the soil and the atmosphere, and carbon that is actually held long term.”<sup>2</sup>



<sup>1</sup> Source: [Grazing and Browsing: How Plants are Affected](#) by Robert K. Lyons and C. Wayne Hanselka – Associate Professor and Professor and Extension Range Specialists, The Texas A&M University System

<sup>2</sup> Source: <https://onpasture.com/2015/11/09/great-grass-farmers-grow-roots/>

August 30, 2021

**Natural Resource Tour of Piceance East-Douglas Herd Management Area (PEDHMA)**


PEDHMA's Appropriate Management Level: **135 – 235 horses**

PEDHMA's (inside and outside) population estimate per BLM Survey Memo 6/10/21: 1,060 horses

**(1,060 horses + 20% annual population increase for 2021 & 2022 = 1,526 horses in 2022)**



Tour participants hearing and observing the results of the range monitoring data conducted 2018 – 2021.



**Close-up view of "grass mortality"**

Years of excess horses in the area caused severe impacts to the rangelands by unmanaged year-long grazing until the root systems could no longer sustain growth and succumb to the pressures.

**Wind and water erosion has removed soil where live plants no longer exist, exposing grass roots.**

Cattle are permitted in the PEDHMA on a seasonal basis.

Horses remain in the PEDHMA year-round, so the grass never gets rest.

# Impacts of Managed Grazing

When not properly managed over the long-term, grazing can have negative impacts to the sustainability of rangeland resources; however, when managed properly grazing can provide ecological benefits compared to no grazing. Research has shown that in arid and semiarid areas, grazing at use levels below 40 percent can have positive impacts to forage plants compared to exclusion of grazing.<sup>[1]</sup>

Research conducted in western Colorado in mountain big sagebrush communities found no significant effects from 40-50 years of grazing exclusion on cover or frequency of grasses, biotic crusts, or bare soil and that grazing exclusion decreased above ground net primary production and biodiversity.<sup>[2]</sup>

In a synthesis of scientific literature on long-term rest in the sagebrush steppe, Davies et al.<sup>[3]</sup> found that long-term rest and properly managed grazing produced few significant differences, and in some situations, negative ecological effects from long-term rest.

September 5, 2021  
Grazing Allotment  
North of Hwy 64  
Rio Blanco County, CO  
Livestock Permitted  
No Horses



Litter from previous years' growth and multiple grass plants hold soil and moisture in place. Livestock are grazed between March and mid-June on a rotational basis. Livestock are managed and remain in each pasture for a maximum of 30 days, allowing for regrowth. Managed grazing in this allotment ensures regrowth opportunities. Managed grazing promotes sustainable rangelands.

[1] Holechek, J.L., T.T. Baker, J. C. Boren, and D. Galt. 2006. Grazing Impacts on Rangeland Vegetation: What We Have Learned. *Rangelands* 28:7-13.

[2] Manier, D.J. and N. T. Hobbs. 2006. Large herbivores influence the composition and diversity of shrub-steppe communities in the Rocky Mountains, USA. *Oecologia* 146: 641. doi:10.1007/s00442-005-0065-9

[3] Davies, K.W., M. Vavra, B. Schultz, and N. Rimbey. 2014. Implications of longer term rest from grazing in the sagebrush steppe. *Journal of Rangeland Applications* 1:14-34.